

# MMR Vaccine (Measles, Mumps, and Rubella): *What You Need to Know*

Many vaccine information statements are available in Spanish and other languages. See [www.immunize.org/vis](http://www.immunize.org/vis)

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite [www.immunize.org/vis](http://www.immunize.org/vis)

## 1. Why get vaccinated?

MMR vaccine can prevent **measles, mumps, and rubella**.

- **MEASLES (M)** causes fever, cough, runny nose, and red, watery eyes, commonly followed by a rash that covers the whole body. It can lead to seizures (often associated with fever), ear infections, diarrhea, and pneumonia. Rarely, measles can cause brain damage or death.
- **MUMPS (M)** causes fever, headache, muscle aches, tiredness, loss of appetite, and swollen and tender salivary glands under the ears. It can lead to deafness, swelling of the brain and/or spinal cord covering, painful swelling of the testicles or ovaries, and, very rarely, death.
- **RUBELLA (R)** causes fever, sore throat, rash, headache, and eye irritation. It can cause arthritis in up to half of teenage and adult women. If a woman gets rubella while she is pregnant, she could have a miscarriage or the baby could be born with serious birth defects.

Most people who are vaccinated with MMR will be protected for life. Vaccines and high rates of vaccination have made these diseases much less common in the United States.

## 2. MMR vaccine

**Children** need 2 doses of MMR vaccine, usually:

- First dose at age 12 through 15 months
- Second dose at age 4 through 6 years

**Infants who will be traveling outside the United States when they are between 6 and 11 months of age** should get a dose of MMR vaccine before travel. These children should still get 2 additional doses at the recommended ages for long-lasting protection.

**Older children, adolescents, and adults** also need 1 or 2 doses of MMR vaccine if they are not already

immune to measles, mumps, and rubella. Your health care provider can help you determine how many doses you need.

A third dose of MMR might be recommended for certain people in mumps outbreak situations.

MMR vaccine may be given at the same time as other vaccines. Children 12 months through 12 years of age might receive MMR vaccine together with varicella vaccine in a single shot, known as MMRV. Your health care provider can give you more information.

## 3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of MMR or MMRV vaccine**, or has any **severe, life-threatening allergies**
- Is **pregnant** or thinks she might be pregnant—pregnant women should not get MMR vaccine
- Has a **weakened immune system**, or has a **parent, brother, or sister with a history of hereditary or congenital immune system problems**
- Has ever had a **condition that makes him or her bruise or bleed easily**
- Has recently had a **blood transfusion or received other blood products**
- Has **tuberculosis**
- Has **gotten any other vaccines in the past 4 weeks**

In some cases, your health care provider may decide to postpone MMR vaccination until a future visit.



U.S. CENTERS FOR DISEASE CONTROL AND PREVENTION

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting MMR vaccine.

Your health care provider can give you more information.

---

#### 4. Risks of a vaccine reaction

---

- Sore arm from the injection or redness where the shot is given, fever, and a mild rash can happen after MMR vaccination.
- Swelling of the glands in the cheeks or neck or temporary pain and stiffness in the joints (mostly in teenage or adult women) sometimes occur after MMR vaccination.
- More serious reactions happen rarely. These can include seizures (often associated with fever) or temporary low platelet count that can cause unusual bleeding or bruising.
- In people with serious immune system problems, this vaccine may cause an infection that may be life-threatening. People with serious immune system problems should not get MMR vaccine.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

---

#### 5. What if there is a serious problem?

---

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your

health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

---

#### 6. The National Vaccine Injury Compensation Program

---

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or call **1-800-338-2382** to learn about the program and about filing a claim.

---

#### 7. How can I learn more?

---

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at [www.fda.gov/vaccines-blood-biologics/vaccines](http://www.fda.gov/vaccines-blood-biologics/vaccines).
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines).

